



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Let $p=2rs$; $q=r^2+s^2$. Then $x=\frac{4r^2s^2}{r+s}$; $y=\frac{(r^2+s^2)^2}{r+s}$.

Let $r=k+l$; $s=k-l$. Then $x=\frac{2(k^2-l^2)^2}{k}$; $y=\frac{2(k^2+l^2)^2}{k}$.

Let $l=\alpha k$. Then $x=2k^3(1-\alpha^2)^2$; $y=2k^3(1+\alpha^2)^2$.

Now $a=p^3=8r^3s^3=8(k^2-l^2)^3=8k^6(1-\alpha^2)^3$, and $b=q^3=(r^2+s^2)^3=8(k^2+l^2)^3=8k^6(1+\alpha^2)^3$, where α and k are integers.

PROBLEMS.

53. Proposed by A. H. BELL, Box 184, Hillsboro, Illinois.

Given $x^2-114xy=\mp 3$ to find the least values of x and y in integers.

54. Proposed by JOSIAH H. DRUMMOND, LL. D., Portland, Maine.

In the expression $2x^2-2ax+b^2$, find two series of values for x in integral terms of a and b .

AVERAGE AND PROBABILITY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

35. Proposed by B. F. FINKEL, A. M., Professor of Mathematics and Physics in Drury College, Springfield, Missouri.


Find the chance that the distance of two points within a square shall not exceed a side of the square. [From *Byerly's Integral Calculus*.]

I. Solution by ALWYN C. SMITH, The University of Colorado, Boulder, Colorado.

a is one side of the square; P and Q the two points; (x, y) the point P with O for origin; and r and ϕ the polar coordinates of Q , with P as origin. Then the favorable cases are

$$4 \int_0^{\frac{1}{2}\pi} \int_0^a \int_0^{a-r\sin\phi} \int_0^{a-r\cos\phi} dx dy r dr d\phi = a^4(\pi - \frac{1}{6}\pi).$$

All the cases $= a^2 \cdot a^2 = a^4$. Therefore, $p = \pi - \frac{1}{6}\pi$.



II. Solution by J. M. COLAW, A. M., Principal of High School, Monterey, Virginia.


Let a = a side of the square $ABCD$, and join A with any point P within the given square. Then as AP represents the distance and direction of the second point from the first, the area of the rectangle $PECF$ represents the number of ways the two points can be taken.

Let $AP = x$, $AH = x'$, and $\angle PAB = \theta$.

When $x' = a \sec \theta$, $PF = a - x \sin \theta$, $PE = a - x \cos \theta$.

\therefore Area $PECF = (a - x \sin \theta)(a - x \cos \theta)$.

Hence the required chance is




$$\begin{aligned} p_1 &= \frac{\int_0^{1\pi} \int_0^a (a - x \sin \theta)(a - x \cos \theta) x dx d\theta}{\int_0^{1\pi} \int_0^{x'} (a - x \sin \theta)(a - x \cos \theta) x dx d\theta} \\ &= \frac{8}{a^4} \int_0^{1\pi} \int_0^a (a - x \sin \theta)(a - x \cos \theta) x dx d\theta \\ &= \frac{8}{a^4} \int_0^{1\pi} (6 - 4 \sin \theta - 4 \cos \theta + 3 \sin \theta \cos \theta) d\theta = \pi - \frac{1}{6}^3. \end{aligned}$$

III. Solution by LEWIS NEIKIRK, Senior in the University of Colorado, Boulder, Colorado.

Take a rectangle $ABCD$, with sides $AB = b$, and $BC = a$, such that a is not greater than b ; and consider the chance that the proposed distance shall exceed b . Let N be the number of favorable cases; then if a be increased infinitesimally, dN will be the number of new cases introduced by placing each point in turn on the differential slice along b while the other one traverses the mixtilinear area DEF .

That is, taking AP equal to x ,



$$\begin{aligned} dN &= 4 \left[\int_{\sqrt{b^2 - a^2}}^b (ax - \frac{a}{2} \sqrt{b^2 - a^2} - \frac{x}{2} \sqrt{b^2 - x^2} \right. \\ &\quad \left. - \frac{b^2}{2} \sin^{-1} \frac{x}{b} + \frac{b^2}{2} \cos^{-1} \frac{a}{b}) dx \right] da \\ &= 2 \left[2ab - ab \sqrt{b^2 - a^2} - \frac{a^3}{3} - \frac{\pi b^3}{2} + b^3 \cos^{-1} \frac{a}{b} \right] da; \text{ and,} \end{aligned}$$

$$N = 2 \left[a^2 b^2 + \frac{b}{3} \sqrt{(b^2 - a^2)^3} - \frac{a^4}{1^2} - \frac{\pi a b^3}{2} + ab^3 \cos^{-1} \frac{a}{b} - b^3 \sqrt{b^2 - a^2} \right] + C.$$


Since $N=0$ when $a=0$, $C=-\frac{4b^4}{3}$; and,

$$N=2[a^2b^2 + \frac{b}{3}\sqrt{(b^2-a^2)^3} - \frac{a^2}{12} - \frac{\pi ab^3}{2} + ab^3\cos^{-1}\frac{a}{b} - b^3\sqrt{b^2-a^2} + \frac{2b^4}{3}].$$

If now $a=b$, $N=b^4(\frac{1}{8}-\pi)$; and the whole number of cases $=b^4$. Hence the chance that the proposed distance shall exceed b is $\frac{1}{8}-\pi$; therefore, the chance that it will not exceed b is $\pi-\frac{1}{8}$.

IV. Solution by LEWIS NEIKIRK, Senior in the University of Colorado, Boulder, Colorado.

Let a be one side of the square, and O the origin. With center O and radius a describe a quadrant. Let P any point within the square (x, y) be one point, and Q be the other point. With center P and radius a describe the circle C . Now Q may be anywhere within the area common to this circle and the square. The favorable cases may then be found by confining Q within the rectangle xy while P traverses the entire square, and then taking four times the result. Hence,



$$p = \frac{4}{a^4} \left\{ \int_0^a \int_0^{\sqrt{a^2-y^2}} xy dx dy + \frac{1}{2} \int_0^a \int_{\sqrt{a^2-y^2}}^a [x\sqrt{a^2-x^2} + y\sqrt{a^2-y^2} + a^2\sin^{-1}\frac{x}{a} - a^2\cos^{-1}\frac{y}{a}] dx dy \right\} = \pi - \frac{1}{8}.$$

V. Solution by G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas-Texas.


This problem affords a splendid test of the correctness of the general value for any convex area as demonstrated in problem 25, page 281, September-October MONTHLY.

Let $AK, AL=p$, $\angle LAB=\theta$, $EF, GH=C$.

For EF , $C=asec\theta$; the limits of p are $asin\theta$ to $acos\theta$.

For GH , $C=psec\theta cosec\theta$; the limits of p are $asin\theta cos\theta$ to $asin\theta$. The limits of θ are 0 to $\frac{1}{2}\pi$.

From problem 25,



$$\Delta = \frac{1}{3a^2} \iint (C^3 - 3a^2C + 2a^3) d\theta dp.$$

$$\begin{aligned} \therefore \Delta &= \frac{8}{3a^4} \int_0^{\frac{1}{2}\pi} \int_{asin\theta cos\theta}^{acos\theta} (p^3 sec^3 cosec^3 \theta - 3a^2 p sec \theta cosec \theta + 2a^3) d\theta dp \\ &\quad + \frac{4}{3a^4} \int_0^{\frac{1}{2}\pi} \int_{asin\theta}^{acos\theta} (a^3 sec^3 \theta - 3a^3 sec \theta + 2a^3) d\theta dp. \end{aligned}$$

$$\Delta = \frac{2}{3} \int_0^{\frac{1}{2}\pi} (\tan\theta \sec^2\theta - 3\sin\theta \cos\theta - 6\tan\theta + 8\sin\theta) d\theta$$

$$+ \frac{4}{3} \int_0^{\frac{1}{2}\pi} (\sec^2\theta - \tan\theta \sec^2\theta - 3 + 3\tan\theta + 2\cos\theta - 2\sin\theta) d\theta.$$

$\therefore \Delta = \frac{1}{8} - \pi. \quad p = 1 - \Delta = \pi - \frac{1}{8} = \text{required chance.}$

PROBLEMS.

44. Proposed by HENRY HEATON, M. Sc., Atlantic, Iowa.

What is the average length of all the chords that may be drawn from one extremity of the major axis of an ellipse if they are drawn at equal angular intervals?

45. Proposed by J. C. WILLIAMS, Boston, Massachusetts.

At the end of the fifth inning the base ball score stands 7 to 9. What is the probability of winning for either team?

46. Proposed by J. A. CALDERHEAD, M. Sc., Professor of Mathematics in Curry University, Pittsburg; Pennsylvania.

Four men starting from random points on the circumference of a circular field and traveling at different rates, take random straight courses across it; find the chance that at least two of them will meet.

MISCELLANEOUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

38. Proposed by S. H. WRIGHT, M. D., A. M., Ph. D., Penn Yan, New York.

In latitude $42^\circ 30'$ north $= \lambda$, at what angle with the horizon will the sun rise, its declination $= 22^\circ$ north $= \delta$?

I. Solution by the PROPOSER.

Let BA be a portion of the equator, $CA = \delta$, a portion of a meridian passing through the sun at C when rising, and describing a small-circle arc CE , parallel with BA , and let BC be a portion of the horizon. Then the angles ECA , and BAC , each $= 90^\circ$, because meridians cut the equator and circles of declination at right angles. Now $CBA = 90^\circ - \lambda$, then $\sin BCA = \sin \lambda \sec \delta = \cos BCE. \therefore BCE = 43^\circ 13' 37'' = \text{required angle.}$

